



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

## A NEW CYNIPID FROM ARIZONA.

BY WILLIAM BEUTENMÜLLER,

NEW YORK CITY.

(PLATE II.)

### ***Aulax chrysothamni*, new species.**

*Male and Female.* — Head rufous, vertex piceous in the female, wholly black in the male, very finely and minutely punctate. Antennæ rufous in the female, piceous in the male. Thorax black, very minutely and evenly punctate, subopaque; parasidal grooves obliterated anteriorly, very fine posteriorly and convergent at the scutellum. Anterior lines from the collar very indistinct and scarcely reaching the middle of the thorax. Median groove from the scutellum wanting. Pleuræ finely striate. Scutellum black, rugose, with two pit-like depressions at the base. Abdomen black in both sexes. Legs rufous. Wings hyaline, with dark brown veins. Length of male 1.50 mm.; of the female 2 to 2.50 mm.

*Gall.* — Polythalamous. White, densely covered with white felt-like substance. Elongated, rounded or club-shaped, enlargements of the terminal twigs of the branches of a species of *Chrysothamnus* (*Bigelovia*), measuring from 15 to 30 mm. in length and 12 to 20 mm. in diameter. Sometimes two or three galls are in a row on the same branch. Internally the gall is white and of a pithy substance.

*Habitat.* — Tucson, Arizona (Gneomar von Krockow).

---

## IS MUTATION A FACTOR IN THE PRODUCTION OF VESTIGIAL WINGS AMONG INSECTS?\*

BY CHARLES T. BRUES,

MILWAUKEE, WIS.


The application by zoölogists of experimental methods to the investigation of the varied problems of evolution has become so general during the past few years, that entomologists have almost entirely neglected to search in any other way for facts bearing on the mutation theory of De Vries.

The wealth of insect species and the constancy of their specific characters render them more available for non-experimental work of this sort than probably any other group of living organisms.

It is with the hope of calling attention to this interesting field that I have been tempted to present the following scattered observations on

---

\*A paper read at the meeting of the Entomological Society of America at Chicago, December 30, 1907.



Galls of *Aulax chrysothamni* n. sp.